



For better
mental health

Understanding post-traumatic stress disorder



'The depth of my sorrow was no surprise to me, but I was totally unprepared for the other reactions – the shakes, nightmares, panic attacks, memory and concentration problems, and the unrelenting exhaustion.'

Bomb attack survivor

'I still have nightmares. They can be triggered by something on television, or the smallest of things, like having the duvet caught round my face. The number of times is lessening, but when they do come I'm learning how to cope. I put the light on, I open the window and say "Yep, look, that was a dream".'

Train crash survivor

'One of my problems resulted from experiencing upsetting flashbacks: people badly injured; the shocked faces of fellow passengers and staff at the hospital; my husband and brother-in-law splattered with blood and mud. They came to me when I was at my most vulnerable. I don't think they will ever go completely, but they are fading and are easier to cope with.'

Train crash survivor

Post-traumatic stress disorder may emerge months or sometimes years after a traumatic experience, affecting people's ability to lead their lives. This booklet describes the causes and the symptoms, and tells you what help is available and how to get it.

?

What is post-traumatic stress disorder?

In recent years, mental health professionals have adopted the term post-traumatic stress disorder (PTSD) to describe a range of psychological symptoms people may experience following a traumatic event, which is outside the normal human experience. The World Health Organisation has defined it as:

'A delayed or protracted response to a stressful event or situation (either short or long-lasting) of an exceptionally threatening or long-lasting nature, which is likely to cause pervasive distress in almost anyone.'

Just hearing news of shattering events, such as the September 11th destruction of New York's World Trade Centre, incidents during the Gulf Wars, the Potter's Bar train crash and the Hillsborough football stadium disaster, can have a lasting impact on individuals. Anybody actually present during a disaster of this nature is likely to become extremely distressed. Every day, people are involved in, or witness, events such as road accidents, muggings, and sexual or physical assaults that cause them deep emotional injury. There is no doubt that the reactions that may follow can seriously hamper and interfere with their lives.

Some survivors have objected to the use of the term disorder, because they see such reactions as an entirely normal and understandable response to abnormal events. But the label recognises that events and social conditions that are beyond our control, and which may fill us with fear or horror, can cause extremely disturbing psychological symptoms.

What are the symptoms?

'He'd almost denied what happened, didn't want to talk about it, didn't want to read any of the articles. Then, one day, a friend mentioned to him that she was worried about her cat, because she had left it that morning trapped up a tree, and it was as though that pushed a button. Dick broke down and shed lots of tears.'

If you have faced a traumatic experience, you may simply feel emotionally numb to begin with, and feelings of distress may not emerge straight away. But sooner or later, you are likely to develop emotional and physical reactions, and changes in behaviour, which may include some of the following:

Reliving aspects of the trauma

- vivid flashbacks (feeling as if the trauma is happening all over again)
- intrusive thoughts and images
- nightmares
- intense distress at real or symbolic reminders of the trauma.



Avoiding memories

- keeping busy
- avoiding situations that remind you of the trauma
- repressing memories (being unable to remember aspects of the event)
- feeling detached, cut off and emotionally numb
- being unable to express affection
- feeling there's no point in planning for the future.

Being easily aroused

- disturbed sleep
- irritability and aggressive behaviour
- lack of concentration
- extreme alertness
- panic response to anything to do with the trauma
- being easily startled.

These responses are all quite normal. Many people find the symptoms will disappear in due course. It's when they last for longer than a month, or when they are very extreme, that PTSD may be diagnosed. Sufferers may also have other symptoms, such as severe anxiety, a phobia or depression. They may develop a dissociative disorder (see *Understanding Dissociative Disorders*, under *Further reading*, on p. 14) and suicidal feelings. There's no time limit on distress, and some survivors may not develop post-traumatic symptoms until many years after the event.



What causes PTSD?

PTSD was first described in relation to the veterans of the Vietnam War in the USA, but the problem has been around for a lot longer under various different names. During the First World War, large numbers of soldiers were said to be suffering from shell shock, soldier's heart or battle fatigue. Rather than admit to the horror of war, and in order to explain the large numbers of men involved, the official line was that these were symptoms of an illness. It has since been estimated that 31 per cent of war veterans suffer from PTSD.

There are important similarities between the psychological stresses associated with war and those that occur in response to major disasters or to personal trauma. Following the sinking of the Herald of Free Enterprise ferry at Zeebrugge, 90 per cent of the passengers and their relatives who were seen for psychological assessment were found to be suffering from PTSD, either alone or in combination with other problems, such as alcohol misuse.

This highlights the fact that people who have not been directly involved in a trauma of this kind may still experience levels of distress as high as those who were. Civilian survivors of war and refugees also develop PTSD, for example. So do people who have lost relatives and friends as a result of some disaster, as well as rescue workers who tried to save them. After the King's Cross underground station fire in 1987, mental health professionals drew up a list of 670 people who were potentially at risk from the psychological effects of the experience. Of that number, only one hundred had been involved directly; the others came from the emergency services, railway officials and the families of victims and survivors. Two years after the Lockerbie air disaster, police, firemen and ambulance workers were still traumatised by what they had witnessed.

Personal trauma, violent crime, sexual or physical assault, road traffic accidents or difficulties during childbirth can all produce PTSD. Adult survivors of childhood sexual abuse may be affected. Children who have themselves been abused, or who have witnessed something terrible, such as a murder, are also prone to the disorder.

Why do some people develop PTSD?

'Every time I hear of any disaster, whether here or abroad, it brings back my own memories, and I can imagine how the families and victims are feeling. My own experience has taught me that you can't change what happened, but that time does make it easier to cope. The memories don't go, but they do fade.'



It's reckoned that about one per cent of the general population is likely to be affected by PTSD at some point. Anyone can develop PTSD following experiences such as those mentioned above, but not everyone does so. Nor does everyone develop it to the same degree. There are a number of possible explanations for this.

Fearing for your life

Events involving loss of life, or where people have been faced by the prospect of dying, may lead to more long-lasting stress responses. A study of Falklands War veterans found that exposure to combat was the most significant factor in predicting who would develop PTSD.

Harmful intentions

Man-made disasters, particularly those involving deliberate acts of violence or terrorism, seem to cause longer-lasting and more painful emotional consequences than natural disasters. The crucial factor may be that it destroys people's trust in others, particularly when it involves someone they depended on.

Conscious memories

People who remain conscious through the experience may be more vulnerable to PTSD because of the horrific memories etched on the mind.

Personal circumstances

Your personal history can make you more prone to PTSD. If a traumatic event triggers memories of an earlier distressing experience, the impact is effectively doubled in intensity. Similarly, if you are already going through emotional problems, you are also much more vulnerable. Your temperament may also have a lot to do with it.

Guilt feelings

Survivors often feel extremely guilty, as though they were responsible for the event, or could have done more to save themselves or others. One study showed that those who blamed themselves in some way for the outcome of the disaster were more at risk of severe and long-term distress.



What's the best way to deal with a traumatic event?

In the aftermath, people often feel numb, dazed and disorientated. Talking about what has happened to them may well be the last thing they want to do. Many survivors have said that what they found most useful, to begin with, was practical advice, information and support with day-to-day tasks.

Expressing your feelings, by talking, may be the best way of coming to terms with the experience. But there are differing views about whether, or not, counselling or some form of psychological de-briefing immediately after the event can prevent serious problems developing later. Everyone will have their own unique responses, and will need to proceed at their own pace. Survivors may turn to friends, relatives and colleagues when they decide they do want to talk about what they've been through. What's important is that there should be an opportunity to talk to someone about their distress, when they're ready to do so. Many of those who do develop PTSD are able to come to terms with the traumatic experience, in this way, within a matter of months.

Stress responses that are bottled up over months or years become deeply ingrained and may cause serious problems. Sometimes, people turn to alcohol or drugs in an attempt to blot out painful feelings and memories. Individuals may remain in a state of extreme tension long after the trauma has passed. They may find themselves avoiding situations, in case they remind them of their trauma, so that life becomes increasingly restricted. In trying to avoid the problem, they are also, in effect, avoiding getting appropriate help. There are various organisations that can provide information about counsellors experienced in treating PTSD. (For more information, see *Useful organisations*, on p. 11.)

It can be very helpful for people to share their experiences with others who have been through something similar. This can be an extremely important step in moving away from isolation, from the role of victim and passive recipient of professional help, towards regaining control of their lives. There are organisations and helplines that specialise in particular groups of people, for

example, soldiers who have seen combat, victims of violent crime or sexual assault, and people who have been tortured or who are refugees. You may find it especially useful to contact an organisation geared to your particular experience (see *Useful organisations*, on p. 11).



What sort of help is available for PTSD?

If you have been suffering from distressing symptoms for months after a traumatic event, you should see your GP, who can refer you for specialist help. There are clinics that focus on PTSD and there are national referral centres. Your GP may also be able to refer you to a local practitioner (a psychologist, psychotherapist or psychiatrist) who has been trained in treating PTSD.

Treatments for PTSD are still being evaluated, but experts agree that for anyone who is still suffering months after the event, the most effective approach is a series of lengthy sessions with a psychologist or other therapist. During these sessions, you will be encouraged to go over the experience in detail.

Many people with PTSD are also extremely depressed. Taking antidepressants may help relieve some of the symptoms and help people to get the best from the psychological treatments. Since these can be addictive, they should be used with caution.

Different types of trauma have different types of impact. Survivors of child abuse may have symptoms that closely resemble PTSD, but they may need much more long-term, intensive help. There is ongoing research being conducted in this area, and specialist programmes have been set up to provide the appropriate therapy for people who were abused as children, victims of political violence, witnesses of murder or those involved in life-threatening situations (see *Useful organisations*, on p. 11). Clinics specialising in treating this kind of problem may provide very structured treatment plans to clients, which involve confronting very painful feelings. They may offer a combination of the following treatments.

Behavioural therapy

This involves going through the trauma, repeatedly, with a psychologist or therapist, so that people re-experience the emotions they felt at the time. It's important to go through the details of the trauma many times until they've dealt with the feelings and feel comfortable and no longer anxious. Some people may not be able to remember the details of the trauma because they were unconscious, or because they have repressed aspects of the experience. They will be encouraged to focus on their fears about the event and any aspects they do remember. Sometimes sessions are recorded on tape, to be listened to later at home. Research suggests that this kind of self-directed remembering can lead to a significant reduction in symptoms.

Behavioural therapy can also help with the problem of avoidance by encouraging people to confront their fears. For example, if you have been avoiding driving since a car crash, the therapist would help you to overcome this, through a step-by-step programme.

Cognitive therapy

Many survivors develop irrational feelings and beliefs as a result of their trauma. Someone who has been raped may start to believe that all men are rapists, for example, or may feel extremely guilty and blame herself. Cognitive therapy involves challenging self-damaging thoughts and beliefs about the event and encourages everyone to replace destructive ideas with a more realistic outlook.

Eye movement desensitisation and reprocessing (EMDR)

This is a relatively new form of therapy, based on behavioural techniques, in which the patient makes rhythmic eye movements while imagining the traumatic event. The eye movements are designed to stimulate the information-processing system in the brain, and the aim of the treatment is to help along the processing of the traumatic events, and speed up readjustment and recovery.



What should friends or relatives do?

'As the shock wore off, I felt under more and more pressure to get on with my life and to behave as though nothing had happened. As a consequence, I became less and less able to do this... It was a great strain trying to balance the need to talk about what had happened, and the consequences, and giving others as little excuse as possible to come out with phrases such as, "Life goes on" and, "You've got to pull yourself together".'

Survivors may need to be given permission to express their feelings about what they've been through. Relatives and friends can help by:

- encouraging them to talk about it
- allowing them to be upset (without necessarily trying to console them)
- getting them to talk to other survivors
- helping them to contact organisations that offer specialist support
- not letting them get into a pattern of avoiding situations that remind them of the trauma.

A traumatic event will have a major impact, not just on those who lived through it, but on their family, friends and colleagues. For sources of support, see *Useful organisations*, opposite.

Useful organisations

ASSIST

(Assistance Support and Self Help in Surviving Trauma)

11 Albert Street, Warwickshire CV21 2QE

helpline: 01788 560 800, tel. 01788 551 919

email: assist@traumatic-stress.freeserve.co.uk

Support, understanding and counselling for people experiencing PTSD, families and carers

British Association for Behavioural and Cognitive Psychotherapies (BABCP)

The Globe Centre, PO Box 9, Accrington BB5 0XB

tel. 01254 875 277, fax: 01254 239 114

email: babcp@babcp.com web: www.babcp.com

Promotes the development of the theory and practice of behavioural and cognitive psychotherapies. Can provide details of accredited therapists. Full directory of psychotherapists available online

British Association for Counselling and Psychotherapy (BACP)

BACP House, 35–37 Albert Street, Rugby CV21 2SG

tel. 0870 443 5252, fax: 0870 443 5161

minicom: 0870 443 5162

email: bacp@bacp.co.uk web: www.bacp.co.uk

See website or send A5 SAE for details of practitioners in your area

Cruse Bereavement Care

Cruse House, 126 Sheen Road, Richmond, Surrey TW9 1UR

helpline: 0870 167 1677

or 0808 808 1677 (for 12 to 18 year olds)

tel. 020 8939 9530, fax: 020 8940 7638

email: helpline@crusebereavementcare.org.uk

or info@crusebereavementcare.org.uk

web: www.crusebereavementcare.org.uk

For all those who have been bereaved

The Compassionate Friends

53 North Street, Bristol BS3 1EN

helpline: 0117 953 9639, tel. 0117 966 5202

email: info@tcf.org.uk web: www.tcf.org.uk

For bereaved parents and their families

First Person Plural

PO Box 2537, Wolverhampton WV4 4ZL

email: fpp@firstpersonplural.org.uk

web: www.firstpersonplural.org.uk

Survivor-led organisation for dissociative survivors and their allies

Ex-Services Mental Welfare Society (Combat Stress)

Tyrwhitt House, Oaklawn Road, Leatherhead KT22 0BX

tel. 01372 841 600

email: contactus@combatstress.org.uk

web: www.combatstress.com

For members of all ranks of the Armed Forces or Merchant Navy. Three treatment centres available

Medical Foundation for the Care of Victims of Torture

96–98 Grafton Road, London NW5 3EJ

tel. 020 7813 7777, fax: 020 7813 0011

email: clinical@torturecare.org.uk

web: www.torturecare.org.uk

Provides survivors of torture with medical treatment, practical assistance and psychotherapeutic support

Refugee Support Centre

47 South Lambeth Road, London SW8 1RH

tel./fax: 020 7820 3606

email: rsctherapy47@hotmail.com

Focuses on the psychological needs of refugees and asylum seekers. People can refer themselves. Support in the language of choice

Traumatic Stress Clinic

73 Charlotte Street, London W1T 4PL

tel. 020 7530 3666, fax: 020 7530 3677

web: www.traumatic-stress-clinic.org.uk

Individual, group or family centred approach by psychologists and psychiatrists. Referral via GP or appropriate agency

Traumatic Stress Service

Maudsley Hospital, 99 Denmark Hill, London SE5 8AZ

tel. 020 7919 2696

National referral centre (via Community Mental Health Teams).

Offers rapid assessment to people suffering from PTSD, or related problems

Victim Support

Cranmer House, 39 Brixton Road, London SW9 6DZ

helpline: 0845 303 0900

Independent charity providing free confidential support to victims of crime

Useful websites

www.ptsd.org.uk

For ex-servicemen and women

www.uktrauma.org.uk

UK Trauma Group

Further reading

- Coping with Post-Trauma Stress* F. Parkinson (Sheldon Press 2000) £6.99
- How to Deal with Bullying at Work* (Mind 2001) £1
- How to Help Someone Who is Suicidal* (Mind 2002) £1
- How to Improve your Mental Wellbeing* (Mind 2002) £1
- How to Look after Yourself* (Mind 2002) £1
- Hurting and Healing: How to overcome the trauma of sexual abuse and rape* G. Wade (Vega 2001) £7.99
- It's my Life Now: Starting over after an abusive relationship or domestic violence* M. Kennedy Dugan, R. R. Hock (Routledge 2000) £11.99
- Making Sense of Antidepressants* (Mind 2002) £3.50
- Making Sense of Cognitive Behaviour Therapy* (Mind 2001) £3.50
- Making Sense of Sleeping Pills* (Mind 2000) £3.50
- The Mind Guide to Managing Stress* (Mind 2003) £1
- The Mind Guide to Physical Activity* (Mind 2001) £1
- The Mind Guide to Relaxation* (Mind 2001) £1
- The Mind Guide to Spiritual Practices* (Mind 2003) £1
- Overcoming Childhood Trauma: A self-help guide using cognitive-behavioural techniques* H. Kennerley (Robinson 2000) £7.99
- Overcoming Traumatic Stress: A self-help guide using cognitive-behavioural techniques* C. Herbert, A. Wetmore (Robinson 1999) £7.99
- A Shattered World: The mental health needs of refugees and newly arrived communities* CVS Consultants and Migrant and Refugee Communities Forum (CVS 2000) £7.50
- Treating Anxiety and Stress: A group psycho-educational approach using brief cognitive behaviour therapy* J. White (John Wiley 2000) £19.99
- Understanding Anxiety* (Mind 2002) £1
- Understanding Depression* (Mind 2003) £1
- Understanding Dissociative Disorders* (Mind 2002) £1
- Understanding Mental Illness* (Mind 2003) £1
- Understanding Talking Treatments* (Mind 2002) £1
- The World is Full of Laughter* D. Sen (Chipmunka Publishing 2003) £10

order form

For a catalogue of publications from Mind, send an A4 SAE to the address below.

If you would like to order any of the titles listed here, please photocopy or tear out these pages, and indicate in the appropriate boxes the number of each title that you require.

Please add 10 per cent for postage and packing, and enclose a cheque for the whole amount, payable to Mind. Return your completed order form together with your cheque to:

Mind Publications

15–19 Broadway

London E15 4BQ

tel. 020 8221 9666

fax: 020 8534 6399

email: publications@mind.org.uk

web: www.mind.org.uk

(Allow 28 days for delivery.)

Please send me the titles marked opposite. I enclose a cheque (including 10 per cent for p&p) payable to Mind for £

Name

Address

Postcode

Tel.

Mind works for a better life for everyone with experience of mental distress

Mind does this by:

- advancing the views, needs and ambitions of people with experience of mental distress
- promoting inclusion through challenging discrimination
- influencing policy through campaigning and education
- inspiring the development of quality services which reflect expressed need and diversity
- achieving equal civil and legal rights through campaigning and education.

The values and principles which underpin Mind's work are:

autonomy, equality, knowledge, participation and respect.

For details of your nearest Mind association and of local services contact Mind's helpline, *MindinfoLine*: **0845 766 0163** Monday to Friday 9.15am to 5.15pm. Speech-impaired or Deaf enquirers can contact us on the same number (if you are using BT Textdirect, add the prefix 18001). For interpretation, *MindinfoLine* has access to 100 languages via Language Line.

Scottish Association for Mental Health tel. 0141 568 7000

Northern Ireland Association for Mental Health tel. 028 9032 8474

This booklet was written by Jane Gorman

First published by Mind 1997. Revised edition © Mind 2003

ISBN 1-876490-27-8

No reproduction without permission

Mind is a registered charity No. 219830

Mind (National Association for Mental Health)

15-19 Broadway

London E15 4BQ

tel: 020 8519 2122

fax: 020 8522 1725

web: www.mind.org.uk



**For better
mental health**